




Meeting the Challenges of Opioids and PAIN

Opioids and Behavioral Health

Tuesday, November 13, 2018



A Provider Toolkit

Meeting the Challenges of Opioids and PAIN:

PATIENT EDUCATION ON PAIN AND OPIOID PRESCRIPTIONS

ADDRESSING OPIOID PRESCRIPTION PRACTICES

IDENTIFYING SAFE AND EFFECTIVE PAIN MANAGEMENT PROTOCOLS

NONPHARMACOLOGIC AND NON-OPIOID PHARMACOTHERAPY ALTERNATIVES



<http://www.stratishealth.org/pip/opioids.html>

Table of Contents

Introduction to the issue.....	2
Shared Decision Making	4
<i>Resources for Shared Decision Making</i>	5
<i>Resources for Patient Education about Pain and Opioids</i>	7
Identifying Opioid Use Disorder or Drug Seeking Behavior	8
<i>Resources for Opioid Use Disorder</i>	9
Prescription Monitoring Programs	10
Effective Screening for Risk Factors	10
Continuing Medical Education and Training Opportunities	12
Non-Pharmacological Alternative Pain Management Therapies	14
<i>Resources for Non-Pharmacologic Interventions</i>	15
Tools for Pharmacists	16
<i>Patient Education</i>	16
<i>Proper Disposal</i>	17
Considerations for the Elderly/Seniors	17
Considerations for Adolescents and Young Adults	18
Safe Storage and Disposal of Prescription Opioids	20
Pet Connection: Opioids Prescribed for Animals	21
Appendix A	23
Resources from Other States and Related Topic Areas:	23
Bibliography	24



Opioids and Behavioral Health


Jessie C Everts, PhD LMFT
Wayside Recovery Center

Objectives

- 1) Identify key components in the intersection of opioid and substance use disorders and mental health issues
- 2) Identify important elements that can have exacerbating effects on both SUD and mental health conditions, and learn how you can be sensitive to these elements in your work.
- 3) Understand how dual diagnosis treatment programs approach healing for mental health and substance use disorders.




Past year SUD and MH diagnoses among US adults



SAMHSA (2015): Results from the 2014 National Survey on Drug Use and Health

COD Treatment

Treatment



2.6 Million Adults with Co-Occurring Serious Mental Illness (SMI) and Substance Use Disorder

Samhsa.gov

Diagnostic Criteria for SUD

- 1: Taking more than intended
- 2: Unsuccessful efforts to decrease use
- 3: Time spent in activities related to use
- 4: Craving
- 5: Failure to fulfill major role obligations
- 6: Interpersonal problems caused by use
- 7: Activities reduced due to use
- 8: Use in situations physically hazardous
- 9: Physical or psychological problems caused by use
- 10: Tolerance
- 11: Withdrawal



Patients diagnosed with opioid use disorder are more frequently diagnosed with...

Relative frequency ratios of select conditions among patients diagnosed vs. not diagnosed with opioid use disorder

Alcoholism	8.4x
Suicidal ideation	6.9x
Binge drinking	5.0x
Bipolar disorder	5.0x
Post traumatic stress disorder	4.2x
Generalized anxiety disorder	2.5x
Depression	2.4x
Insomnia	2.0x

Alcoholism is **8.4 times** more frequently diagnosed among patients also diagnosed with opioid use disorder versus patients not diagnosed with opioid use disorder.



Based on 3.1 million patients diagnosed with opioid use disorder and at least one other condition in 2014-2016. Take the guesswork out of healthcare on amino.com

© 2017 Amino Inc.

Opioids and Mental Health

HIGHER OPIOID PRESCRIPTION RATES FOR THOSE WITH MENTAL HEALTH DISORDERS



Only 5% of **Americans without mental health disorders** are prescription opioid users.



Over 18% of **Americans with mental health disorders** are prescription opioid users.



Americans with mental health disorders make up 16% of the U.S. population...



but are prescribed over 51% of the **115 million opioid prescriptions** given in the U.S. in 2016.

Data from: "Prescription Opioid Use Among Adults with Mental Health Disorders in the United States" Journal of the American Board of Family Medicine, July 6, 2017



Opioids – Strategies

- **Emergency Response** (Tertiary Prevention) – Naloxone, Good Samaritan Laws, Syringe Exchange, transitions of care, discharge planning, fentanyl alerts, infectious disease control
- **Intervention & Treatment** (Secondary Prevention) – Screening, early identification, SBIRT Services, Medication Assisted Therapy, chemical health treatment, OB and infant care for NAS, safe storage, safe disposal
- **Primary Prevention & Public Health** (Primary Prevention) – Prescribing practices, safe use of prescriptions, control supply, prevent diversion, reduce marketing, enrollment and use of PMP, prevention of ACEs, adolescent risk reduction, pain management, addressing trauma, integrating care, protective factors, community resiliency, culture as prevention



Opioids – Long Term Goals

Statewide goals – The overarching, long terms goals are to:


- Reduce opioid overdose death
- Reduce non-fatal opioid overdose
- Reduce disparity in opioid use and overdose death within Native American, African American, and LGBTQ communities
- Improve prescribing practices
- Diagnose and treat substance use disorder
- Increase investment, intention and financial support for identified high-need communities
- Increase the conditions that support health (e.g. mental, medical, dental, sexual, chemical)



Social Determinants

Determinants of Health

Factors that contribute to the social patterning of health, disease, and illness



Minnesota
Department of
Health



ASAM Six-Dimension Assessment

AT A GLANCE: THE SIX DIMENSIONS OF MULTIDIMENSIONAL ASSESSMENT


ASAM's criteria uses six dimensions to create a holistic, biopsychosocial assessment of an individual to be used for service planning and treatment across all services and levels of care. The six dimensions are:

1	DIMENSION 1	Acute Intoxication and/or Withdrawal Potential Exploring an individual's past and current experiences of substance use and withdrawal
2	DIMENSION 2	Biomedical Conditions and Complications Exploring an individual's health history and current physical condition
3	DIMENSION 3	Emotional, Behavioral, or Cognitive Conditions and Complications Exploring an individual's thoughts, emotions, and mental health issues
4	DIMENSION 4	Readiness to Change Exploring an individual's readiness and interest in changing
5	DIMENSION 5	Relapse, Continued Use, or Continued Problem Potential Exploring an individual's unique relationship with relapse or continued use or problems
6	DIMENSION 6	Recovery/Living Environment Exploring an individual's recovery or living situation, and the surrounding people, places, and things




ASAM: American Society of Addiction Medicine

REFLECTING A CONTINUUM OF CARE



Co-occurring Care at Wayside



Trauma-informed Care in a nutshell

TRAUMA-INFORMED CARE: GUIDING VALUES **"HEALING HAPPENS IN RELATIONSHIP"**



What happens in treatment?

Most common modalities:

- Cognitive Behavior Therapy (CBT)
- Illness Management & Recovery (IMR)
- Dialectical Behavior Therapy (DBT)
- Family Therapy
- Wellness/Life Skills groups
- Use/Relapse Prevention skills groups
- 12-Step Facilitation
- Seeking Safety/Trauma therapies
- Mental Health therapy
- Culturally specific programs
- Medication-assisted treatment (MAT)
- Case Management/Care Coordination
- Peer Recovery Services (PRS)



What gets in the way of treatment effectiveness

Client factors and Provider factors!

- Shame
- Stigma
- Separation or threat of separation from children
- Legal consequences
- Financial barriers
- Fragmented/siloed systems
- Abstinence-only/zero-tolerance philosophies
- Medication adherence
- Lack of social support
- Symptoms



Why focus on Shame?

Shame gets in our way when we want to make a change.

How **Shame** Changes
Your **Client's Brain**

NICABM (2018)



1. Activates default
mode network



2. Activates pain
system

WHAT'S WRONG WITH YOU

WHAT OTHER PEOPLE
THINK ABOUT YOU

WHAT'S WRONG WITH
OTHER PEOPLE

Why focus on Stigma?

Stigma also gets in our way when we want to make a change.

- **Stigma** is real or perceived judgment from others, and is a major deterrent to seeking treatment.
- 3 out of 4 people with mental illness report feeling stigmatized.
- Misconception about mental illness is that it is under the person's control.
- Message: "You did this to yourself, fix it yourself" from doctors, media, words we use, attitudes.



Ways to Corrode Stigma

We have to make it okay to talk about, to seek treatment, and restore dignity (sense of worth) for those suffering from SUD/mental illness.

- Person-first language
- Telling the stories
- EMPATHY (is a skill)
- Advocate: Talk, Listen, Learn



Language Matters – *Person First*

Think about how your words reflect your attitudes and may trigger shame for clients:

Stigmatizing Language	Preferred Language
Addict/User	Person with SUD
Clean	Abstinent
Dirty	Actively using
Relapse/Slip	Resumed/Experienced a recurrence of use
Former addict/alcoholic	Person in Recovery
Clean/Dirty UA or screen	Negative/Positive UA or screen
NAS baby	Infant with NAS
Failed treatment/episode	Prior treatment episode





Ways to Corrode Shame and Stigma

- Recognize things providers say that inadvertently trigger shame (which can be debilitating in making changes)
- Provide a safe space that embraces vulnerability
- Encourage people to talk about their stories by being non-judgmental and asking questions to help them identify their own motivators

Helping someone connect to SUD Treatment

- **Talk with patients about their substance use and recognize the six different dimensions of risk**
- **Be able to answer some questions about the treatment experience**
- **Do some connecting – know where to send someone for a Rule 25/Comprehensive Assessment**
- **Follow up – motivation changes day by day**



Helping someone connect to SUD Treatment

Motivation for self-initiation

Use Motivational Interviewing techniques:

- Express empathy through reflective listening.
- Explore discrepancies between clients' goals or values and their current behavior.
- Avoid argument and direct confrontation.
- Adjust to client resistance rather than opposing it directly.
- Support self-efficacy and optimism.



Takeaways

SUD Treatment is hard work!

There are a lot of barriers to getting into and being successful in SUD treatment – some barriers can be minimized by:

- Working together as providers
- Recognizing the impact of shame and stigma
- Adapting our view of people with SUD
- Being aware of resources and supports

